Recap of Longitudinal Methods



- Repeated measurements over time.
- Unique because they involve correlation within subjects (not independent).

- Repeated measurements over time.
- Unique because they involve correlation within subjects (not independent).
- Require special techniques to analyze them!

- Repeated measurements over time.
- Unique because they involve correlation within subjects (not independent).
- Require special techniques to analyze them!
- ▶ In practice we need to consider wide and long formatted data.

- Repeated measurements over time.
- Unique because they involve correlation within subjects (not independent).
- Require special techniques to analyze them!
- ▶ In practice we need to consider **wide** and **long** formatted data.
- Be mindful of **time scale** and whether or not the data are **balanced**.

Extension of GLMs to longitudinal data.

- Extension of GLMs to longitudinal data.
- Specify a link function, linear predictor, variance function, and correlation pattern.

- Extension of GLMs to longitudinal data.
- Specify a link function, linear predictor, variance function, and correlation pattern.
- ▶ Represent models for **population-level** effects.

- Extension of GLMs to longitudinal data.
- Specify a link function, linear predictor, variance function, and correlation pattern.
- ► Represent models for **population-level** effects.
- Using M-estimation theory, can be estimated through GEEs.

- Extension of GLMs to longitudinal data.
- Specify a link function, linear predictor, variance function, and correlation pattern.
- ► Represent models for **population-level** effects.
- ▶ Using M-estimation theory, can be estimated through **GEEs**.
- In the continuous case, there is a likelihood alternative.

- Extension of GLMs to longitudinal data.
- Specify a link function, linear predictor, variance function, and correlation pattern.
- ► Represent models for **population-level** effects.
- Using M-estimation theory, can be estimated through GEEs.
- In the continuous case, there is a likelihood alternative.
- Uses nlme::gls() or geepack::geeglm() in R.

> Allow for individual-level effects to be estimated in addition to population-level.

Allow for individual-level effects to be estimated in addition to population-level.
Accounts for correlation by adding random effects terms.

- > Allow for individual-level effects to be estimated in addition to population-level.
- Accounts for correlation by adding random effects terms.
- Explicitly models the full structure of the process, including distributional assumptions.

- > Allow for individual-level effects to be estimated in addition to population-level.
- Accounts for correlation by adding random effects terms.
- Explicitly models the full structure of the process, including distributional assumptions.
- Requires likelihood theory to fit.

- > Allow for individual-level effects to be estimated in addition to population-level.
- Accounts for correlation by adding random effects terms.
- Explicitly models the full structure of the process, including distributional assumptions.
- Requires likelihood theory to fit.
- Uses nlme::lme() in R.

▶ Models the process of **categorical** longitudinal data, as a **stochastic process**.

Models the process of categorical longitudinal data, as a stochastic process.
 Simplified by making an *r*-th order Markov assumption or time homogeneity.

- ▶ Models the process of **categorical** longitudinal data, as a **stochastic process**.
- Simplified by making an *r*-th order **Markov** assumption or time homogeneity.
- Likelihood procedure without covariates, using transition counts.

- ▶ Models the process of **categorical** longitudinal data, as a **stochastic process**.
- Simplified by making an *r*-th order **Markov** assumption or time homogeneity.
- Likelihood procedure without covariates, using transition counts.
- Conditional likelihood procedure with covariates, using standard GLM.

- Models the process of categorical longitudinal data, as a stochastic process.
- Simplified by making an *r*-th order **Markov** assumption or time homogeneity.
- Likelihood procedure without covariates, using transition counts.
- Conditional likelihood procedure with covariates, using standard GLM.
- Uses table() or glm() in R.

► Longitudinal data are often subject to severe missingness.

Longitudinal data are often subject to severe missingness.
 Missingness can be MCAR, MAR, or NMAR.

- Longitudinal data are often subject to severe missingness.
- Missingness can be MCAR, MAR, or NMAR.
- Missingness is easier to handle when it is monotone (e.g., with dropout).

- Longitudinal data are often subject to severe missingness.
- Missingness can be MCAR, MAR, or NMAR.
- Missingness is easier to handle when it is monotone (e.g., with dropout).
- Can use weighting techniques or imputation to account for it.

- Longitudinal data are often subject to severe missingness.
- Missingness can be MCAR, MAR, or NMAR.
- Missingness is easier to handle when it is monotone (e.g., with dropout).
- Can use weighting techniques or imputation to account for it.
- Uses mice::mice() or glm() in R.

What comes Next?

The analysis of **survival data**.